

IGR J18483–0311: a new intermediate supergiant fast X-ray transient^{*}

F. Rahoui^{1,2} and S. Chaty¹

¹ Laboratoire AIM, CEA/DSM - CNRS - Université Paris Diderot, Irfu/Service d’Astrophysique, Bât. 709, CEA/Saclay, F-91191 Gif-sur-Yvette, France, e-mail: frahoui, chaty@cea.fr

² European Southern Observatory, Alonso de Córdova 3107, Vitacura, Santiago de Chile

Received; accepted

ABSTRACT

Context. IGR J18483–0311 is a high-mass X-ray binary recently discovered by *INTEGRAL*. Its periodic fast X-ray transient activity and its position in the Corbet diagram - although ambiguous - led to the conclusion that the source was a likely Be/X-ray binary (BeXB), even if a supergiant fast X-ray transient (SFXT) nature could not be excluded.

Aims. We aimed at identifying the companion star of IGR J18483–0311 to discriminate between the BeXB and the SFXT nature of the source.

Methods. Optical and near-infrared photometry, as well as near-infrared spectroscopy of the companion star were performed to identify its spectral type. We also assembled and fitted its broad-band spectral energy distribution to derive its physical parameters.

Results. We show that the companion star of IGR J18483–0311 is an early-B supergiant, likely a B0.5Ia, and that its distance is about 3–4 kpc.

Conclusions. The early-B supergiant nature of its companion star, as well as its fast X-ray transient activity point towards an SFXT nature of IGR J18483–0311. Nevertheless, the long duration and the periodicity of its outbursts, as well as its high level of quiescence, are consistent with IGR J18483–0311 being an intermediate SFXT, in between classical supergiant X-ray binaries (SGXBs) characterised by small and circular orbits, and classical SFXTs with large and eccentric orbits.

Key words. Infrared: stars – X-rays: binaries, individuals: IGR J18483–0311 – binaries: general – supergiants – Stars: fundamental parameters

1. Introduction

High-mass X-ray binaries (HMXBs) are X-ray sources for which high-energy emission stems from accretion onto a compact object of material coming from a massive companion star. Until recently, the majority of known HMXBs were Be/X-ray binaries (BeXBs), and just a few were supergiant X-ray binaries (SGXBs). The launch of the *INTErnational Gamma-Ray Astrophysics Laboratory* (*INTEGRAL*, Winkler et al. 2003) in October 2002 completely changed the situation, as many more HMXBs whose companion stars are supergiants were discovered during the monitoring of the Galactic centre and the Galactic plane using the onboard IBIS/ISGRI instruments (Ubertini et al. 2003; Lebrun et al. 2003). Most of these sources are reported in Bird et al. (2007) and Bodaghee et al. (2007), and their studies reveal two main features which were not present on previously known SGXBs: first, many of them exhibit a considerable intrinsic absorption, far above the interstellar level; second, some of these new sources revealed a transitory nature and occasionally exhibit a fast X-ray transient activity lasting a few hours. It then appears that the *INTEGRAL* supergiant HMXBs can be classified in two classes: one class of considerably obscured persistent sources (see e.g. Chaty & Rahoui 2007) and another of transitory sources called supergiant fast

X-ray transients (SFXTs, Negueruela et al. 2006).

IGR J18483–0311 was discovered during observations performed with *INTEGRAL* in 2003 April 23–28 (Chernyakova et al. 2003), and it was found to have average fluxes of about 10 mCrab and 5 mCrab in the 15–40 keV and 40–100 keV bands, respectively. Molkov et al. (2004) also detected the source in 2003 March–May during a survey of the Sagittarius arm tangent with *INTEGRAL*, and gave average fluxes of about 4.3 mCrab and 3.9 mCrab in the 18–60 keV and the 60–120 keV bands, respectively. Moreover, analysing archival data from the *Rossi X-ray Timing Explorer* (*RXTE*) All-Sky Monitor (ASM), Levine & Corbet (2006) reported a 18.55 days periodicity of the lightcurve of IGR J18483–0311, likely corresponding to its orbital period.

With archival data from observations performed with *INTEGRAL* from 2003 May to 2006 April, Sguera et al. (2007) found five new outbursts from IGR J18483–0311 whose activity lasted no more than a few days, and characterised by fast flares lasting a few hours. They fitted its 3–50 keV spectrum with an absorbed cut-off power law, and derived $N_{\text{H}} \sim 9 \times 10^{22} \text{ cm}^{-2}$ (well above the galactic column density in the line of sight of about $1.6 \times 10^{22} \text{ cm}^{-2}$), $\Gamma \sim 1.4$, and $E_{\text{c}} \sim 22 \text{ keV}$. They also showed that all the outbursts are well-fitted by an absorbed bremsstrahlung whose parameters are $N_{\text{H}} \sim 7.5 \times 10^{22} \text{ cm}^{-2}$ and $kT \sim 21.5 \text{ keV}$. They reported a periodicity in the long term ISGRI light curve of about 18.52 days - confirming the result of Levine & Corbet (2006) - and argued it is most likely an orbital

Send offprint requests to: F. Rahoui

^{*} Based on observations carried out at the European Southern Observatory under the Target of Opportunity programme ID 078.D-0268.

period. They also derived a periodicity of about 21.05 s in the JEM-X light curve of the first outburst, which is likely a neutron star pulse period. Finally, from archival *Swift* observations, the authors obtained a very accurate position of IGR J18483–0311, which allowed them to pinpoint its USNO B1.0 0868–0478906 and 2MASS J18481720–0310168 counterparts. They found that the magnitudes were typical of an absorbed massive late O/early B star which, along with the position of the source in the Corbet diagram as well as the periodicity of its high-energy activity, led them to conclude that IGR J18483–0311 was likely a BeXB, without excluding the possibility of an SFXT nature.

Chaty et al. (2008) assembled the broad-band spectral energy distribution (SED) of the companion star of IGR J18483–0311 from 0.7 μm to 8 μm , with near-infrared (NIR) photometric observations performed at the ESO/NTT using the SofI instrument, as well as archival optical data from the USNO-A.2 catalogue, and mid-infrared (MIR) data from the GLIMPSE survey. They fitted it with a combination of two absorbed black bodies and showed that the best fit was consistent with the companion star being a heavily absorbed B star, confirming the HMXB nature of IGR J18483–0311. At last, Masetti et al. (2008) performed optical spectroscopy of the companion star of IGR J18483–0311, and concluded that it was an O/B giant star because of the large equivalent width of the H_α emission line.

In this paper, we report optical and NIR observations of IGR J18483–0311 performed in service mode at the ESO/NTT through our Target of Opportunity program ID 078.D-0268 (P.I. S. Chaty). These observations aimed at constraining the nature of its companion star and of the binary system. In Sect. 2, we describe the optical/NIR photometric and spectroscopic observations, as well as their reduction. In Sect. 3, we present the results and show the broad-band SED of the companion star of IGR J18483–0311. We finally discuss the outcomes in Sect. 4.

2. Observations and data analysis

2.1. Optical and NIR photometry

In 2007 March 24, we performed optical photometry of IGR J18483–0311 in *U*, *B*, *V*, *R*, and *I* bands with the imager SUSI2 installed at ESO/NTT. We used the large field of view (5'5×5'5), and the plate scale was 0'16 per pixel. The exposure time was 60 s in all filters for each exposure. We also observed four photometric standard stars from the optical standard catalogue of Landolt (1992): PG1657+056, PG1657+056A, PG1657+056B, and PG1657+056C.

NIR photometry was carried out in *J*, *H*, and *Ks* bands with the spectro-imager SofI at ESO/NTT. We used the large field of view (4'92×4'92), and the plate scale was 0'288 per pixel. In each filter, the observations were obtained by repeating a set of images with 9 different 30'' offset positions to subtract the sky emission, and the exposure time was 1.2 s for each position. We also observed three photometric standard stars from the faint NIR standard star catalogue of Persson et al. (1998): sj9116, sj9150, and sj9155.

The photometric data were reduced using *IRAF* by performing flatfielding and bad pixels correction, as well as bias subtraction in the optical bands, and crosstalk correction and sky subtraction in the NIR. After the reduction, all the images were aligned and combined. We then carried out aperture photometry on the optical images. In the NIR, we used the software *StarFinder*, part of the *Scisoft* package from ESO, well-suited

for point-source photometry in crowded fields. The instrumental magnitudes were then transformed to apparent magnitudes using the zero-point, the extinction coefficient, and the color term corrections computed by linear fits using the standard stars. The optical and NIR magnitudes are listed in Table 1. It is worth noting that our optical magnitudes are about 1 magnitude lower than those given in the USNO catalogue, this difference probably due to the strong uncertainty (≥ 1 magnitude) of the USNO magnitudes for very faint sources (≥ 17 in the *B* band).

2.2. NIR spectroscopy

In 2007 March 18-19, we performed NIR low-resolution ($R \sim 1000$) spectroscopy with SofI between 0.9 μm and 2.5 μm , using the blue and red grisms and the 1'' slit. Twelve spectra were taken in each grism, half of them on the source and the other half with an offset of 30'' and allowing for sky emission subtraction. The exposure time of each spectrum was set to 60 s. We also observed in the same conditions HIP96265, a G3V telluric standard star, for telluric features correction.

The NIR spectra were reduced with *IRAF* by performing crosstalk correction, flatfielding, sky subtraction, and bad pixels correction. The target spectra were then extracted, wavelength calibrated by extraction of a xenon arc with the same setup, and finally combined. The telluric features were corrected by dividing the final target spectrum by the telluric standard. To avoid line contamination by the standard star, we finally multiplied the corrected spectrum by a solar spectrum provided by the National Solar Observatory (NSO), and modified using an *IRAF* routine developed by Maiolino et al. (1996). The routine corrects the solar spectrum for airmass, heliocentric, and rotational velocity to match the standard star properties, and smoothes the spectrum to rebin it to the instrument resolution. The final spectrum of the companion star of IGR J18483–0311 - obtained by combining the blue and red grisms - is displayed in Fig 1.

3. Results


3.1. Spectral classification

To identify the spectral type of the companion star of IGR J18483–0311, we compared our spectrum with spectra from the *H* and *K* bands hot stars classifications given in Hanson et al. (1996, 2005) and the *K* band spectroscopic classification given in Clark & Steele (2000). In the spectrum displayed in Fig 1, we marked all the detected features (listed in Table 2), most of them due to $H\text{I}$ and HeI . In the *H* band spectrum, we also detected the $\text{Pa}\beta$ (5–3) feature at 1.2822 μm and the O I feature at 1.3170 μm .

The presence of the $\text{Br}\gamma$ (7–4) line in absorption at 2.1668 μm and of several HeI lines is typical of a normal O9-B3 massive star (Clark & Steele 2000). This therefore excludes the system to be a BeXB. Moreover, the presence of the HeI line at 2.0580 μm in emission and the HeI line at 2.1121 μm in absorption is typical of early B0-B2 supergiants. Comparing the relative strengths of the HeI lines at 1.6996 μm , 2.0580 and 2.1121 μm , and of the $H\text{I}$ lines at 1.6819 μm , 1.7362 μm , and 2.1668 μm , with the relative strengths of the same lines in Hanson et al. (1996, 2005), we find that the star is likely a B0.5Ia supergiant.

Table 1. Optical and NIR magnitudes (Mag) of IGR J18483–0311 obtained with SUSI2 and SofI. We also give the zero point Z_p in each filter.

Filters	<i>U</i>	<i>B</i>	<i>V</i>	<i>R</i>	<i>I</i>	<i>J</i>	<i>H</i>	<i>Ks</i>
Z_p	23.279 ± 0.023	24.837 ± 0.030	24.423 ± 0.018	24.480 ± 0.013	22.707 ± 0.061	23.059 ± 0.020	22.814 ± 0.013	22.211 ± 0.019
Mag	> 23.702	> 25.162	21.884 ± 0.313	17.888 ± 0.041	14.382 ± 0.061	10.840 ± 0.028	9.376 ± 0.018	8.472 ± 0.023

**Fig. 1.** 1.12 μm to 2.3 μm low-resolution spectrum of the companion star of IGR 18483–0311, obtained with the blue and red grisms of SofI.**Table 2.** Spectroscopic features we detected in the blue and red spectra of the companion star of IGR J18483–0311. We also give their laboratory and fitted wavelengths in μm (λ and λ_{fit} , respectively), their equivalent width ($\tilde{W}$) and their full-width at half-maximum ($FWHM$), in $\text{\AA}$.

Identification	λ	λ_{fit}	$\tilde{W}$	$FWHM$
Pa (5-3)	1.2822	1.2820	1.013	8.351
He I	1.2976	1.2975	0.765	20.97
O I	1.3165	1.3170	1.700	17.4
Br (23-4)	1.5043	1.5050	0.949	22.53
Br (19-4)	1.5265	1.5264	1.018	41.25
Br (15-4)	1.5705	1.5698	0.856	23.40
Br (13-4)	1.6114	1.6111	1.049	37.56
Br (12-4)	1.6412	1.6405	0.641	25.94
Br (11-4)	1.6811	1.6819	0.723	27.25
He I	1.7007	1.6996	1.053	32.72
Br (10-4)	1.7367	1.7362	0.813	33.26
He I	2.0587	2.0580	−5.794	32.16
He I	2.1126	2.1121	1.168	30.67
Br (7-4)	2.1661	2.1668	1.151	23.17

3.2. Spectral energy distribution

To assemble the broad-band SED of IGR J18483–0311, we added its archival MIR fluxes from the *Spitzer*’s Galactic Legacy Infrared Mid-Plane Survey Extraordinaire (GLIMPSE, Benjamin et al. 2003) of the Galactic plane ($|b| \leq 1^\circ$ and be-

tween $l=10^\circ$ and $l=65^\circ$, on both sides of the Galactic centre) performed with the *Spitzer Space Telescope* with the IRAC camera in four bands, $3.60 \pm 0.38 \mu\text{m}$, $4.50 \pm 0.51 \mu\text{m}$, $5.80 \pm 0.73 \mu\text{m}$, and $8.00 \pm 1.43 \mu\text{m}$. The GLIMPSE fluxes of the companion star of IGR J18483–0311 are listed in Chaty et al. (2008). We then fitted the SED (using a χ^2 minimisation) with an absorbed black body representing the companion star emission, following Rahoui et al. (2008). We added to the flux uncertainties a 5% systematic error in each NIR band, and a 2% systematic error in each IRAC band, as given in the SofI¹ and the IRAC² manuals. The best-fitted SED and parameters are displayed in Fig 2 and listed in Table 3, respectively.

3.3. Distance

The best-fitted temperature of the companion star of IGR 18483–0311 ($T_* = 24600 \text{ K}$) is consistent with the star being a B0.5Ia supergiant, following the temperature calibration given in Searle et al. (2008). In their paper, they also give the expected unabsorbed absolute magnitude of a B0.5Ia supergiant star, $M_v = -6.5$, as well as its expected radius, $R_e = 33.8 R_\odot$. Using these two parameters, as well as A_v and R_*/D_* that we derived from our fit, we can assess the distance of the source with one of the following relations: $D_* = 10^{0.2(m_v - A_v - M_v + 5)}$ (in pc), or $D_* = \frac{R_e}{R_*/D_*}$ (in

¹ <http://www.la.eso.org/docs/LSO-MAN-ESO-40100-0004/LSO-MAN-ESO-40100-0004.pdf>

² <http://ssc.spitzer.caltech.edu/documents/som/som8.0.irac.pdf>




Fig. 2. Broad-band SED of IGR J18483–0311. Assembled with optical, near-, and mid-infrared data from SUSI2, SofI, and IRAC, respectively.

Table 3. Best-fitted parameters of the SED of IGR J18483–0311. We give the total galactic absorption in the line of sight due to H I ($A_{\text{H I}}$) and H₂ (A_{H_2}), the absorption of the source in the X-ray domain A_{x} , and then the parameters: the optical intrinsic absorption A_{v} , the temperature T_* and the R_*/D_* radius-to-distance ratio of the star. The absorptions are given in magnitudes.

$A_{\text{H I}}$	A_{H_2}	A_{x}	A_{v}	$T_*[\Delta T_*](\text{K})$	R_*/D_*
7.9	9.6	48.1	$15.7^{+0.2}_{-0.3}$	24600[17500 – 36800]	$1.94^{+0.39}_{-0.41} \times 10^{-10}$

in $R_\odot$), where m_v is the optical apparent magnitude of the source. Both relations give $D_* = 3.44$ kpc and $D_* = 3.93$ kpc, respectively. Both values are consistent within the uncertainties and we therefore conclude that the distance of IGR J18483–0311 is about 3–4 kpc.

4. Discussion and conclusions

The identification of its companion star as a B0.5Ia supergiant, as well as its fast X-ray transient activity, are consistent with IGR J18483–0311 being an SFXT. Nevertheless, as already pointed out in Sguera et al. (2007), the IGR J18483–0311 X-ray behaviour appears unusual. Indeed, whereas outbursts last a few hours and $L_{\text{max}}/L_{\text{min}} \sim 10^4$ for typical SFXTs, the outbursts of IGR J18483–0311 last a few days and its $L_{\text{max}}/L_{\text{min}}$ ratio is $\sim 10^3$ (meaning that its quiescence is higher). Moreover, Levine & Corbet (2006) and Sguera et al. (2007) reported a 18.52 days orbital period, which is quite low compared to what is expected for classical SFXTs, with large and eccentric orbits.

Recently, several authors pointed out the importance of clumpy winds to explain the SFXT behaviour (in’t Zand 2005; Leyder et al. 2007; Walter & Zurita Heras 2007; Negueruela et al. 2008). They argue that the flares are created by the interaction of the compact object with dense clumps (created in the stellar wind of the companion star), the frequency of the flares depending mainly on the geometry of the system. On the contrary, the quiescent emission would be due to the accretion onto the compact object of diluted inter-clumps medium, which would explain the very low quiescence exhibited by classical SFXTs. Moreover, by classifying twelve SFXTs in

function of the duration and the frequency of their outbursts, as well as their $L_{\text{max}}/L_{\text{min}}$ ratio, Walter & Zurita Heras (2007) showed the existence of intermediate SFXTs, characterised by a lower ratio and longer flares. Finally, Negueruela et al. (2008) proposed a general scheme to explain the emission of both SGXBs and SFXTs. The authors argued about the existence of a zone around the companion supergiant star (radius $\leq 2R_*$), in which the clump density is very high, and another one (radius $\geq 3R_*$) in which it is low. In function of the orbital parameters of the compact object, the system could be a classical SGXB (small and circular orbit inside the zone of high clump density), a classical SFXT (large and eccentric orbit), or an intermediate SFXT (small orbits, circular or eccentric) with possible periodic outbursts (see also Sidoli et al. 2007, for an alternative model).

Considering the longer duration of its flares as well as their 18.52 days periodicity, and its lower $L_{\text{max}}/L_{\text{min}}$ ratio, IGR J18483–0311 is likely an intermediate SFXT. Moreover, to figure out - in the framework of the model proposed by Negueruela et al. (2008) - whether its orbit is circular or elliptic, we can first use the Kepler’s third law for a B0.5Ia supergiant mass and radius $M_* = 33M_\odot$ and $R_* = 33.8R_\odot$ (Searle et al. 2008), and a neutron star mass of $1.4M_\odot$ to derive the semi-major axis and we find $a \sim 2.83R_*$. Then, assuming that IGR J18483–0311 only gets into activity when orbiting within the zone of high clump density, we find that an eccentricity $0.43 \leq e \leq 0.68$ is needed to explain the average three days bursting activity reported in Sguera et al. (2007). We then conclude that IGR J18483–0311 is an intermediate SFXT with a small and eccentric orbit.

Acknowledgements. We are grateful to Jérôme Rodriguez for his useful website in which all *INTEGRAL* sources are referenced (<http://isdc.unige.ch/~rodrigue/html/igrsources.html>). F.R. acknowledges the CNRS/INSU for the funding of the third year of his ESO/CEA PhD studentship. This research has made use of NASA’s Astrophysics Data System, of the SIMBAD and VizieR databases operated at CDS, Strasbourg, France, of products from the US Naval Observatory catalogues, of products from the Two Micron All Sky Survey as well as products from the Galactic Legacy Infrared Mid-Plane Survey Extraordinaire, which is a *Spitzer Space Telescope* Legacy Science Program.

References

- Benjamin, R. A., Churchwell, E., Babler, B. L., et al. 2003, *PASP*, 115, 953
- Bird, A. J., Malizia, A., Bazzano, A., et al. 2007, *ApJS*, 170, 175
- Bodaghee, A., Courvoisier, T. J.-L., Rodriguez, J., et al. 2007, *A&A*, 467, 585
- Chaty, S. & Rahoui, F. 2007, *Proceeding of the Vith INTEGRAL Workshop, "The Obscured Universe"*, Space Research Institute, Moscow, Russia, 2006, ESA’s Publications Division: Special Publication SP-622
- Chaty, S., Rahoui, F., Foellmi, C., et al. 2008, *A&A*, 484, 783
- Chernyakova, M., Lutovinov, A., Capitanio, F., Lund, N., & Gehrels, N. 2003, *The Astronomer’s Telegram*, 157, 1
- Clark, J. S. & Steele, I. A. 2000, *A&AS*, 141, 65
- Hanson, M. M., Conti, P. S., & Rieke, M. J. 1996, *ApJS*, 107, 281
- Hanson, M. M., Kudritzki, R.-P., Kenworthy, M. A., Puls, J., & Tokunaga, A. T. 2005, *ApJS*, 161, 154
- in’t Zand, J. J. M. 2005, *A&A*, 441, L1
- Landolt, A. U. 1992, *AJ*, 104, 340
- Lebrun, F., Leray, J. P., Lavocat, P., et al. 2003, *A&A*, 411, L141
- Levine, A. M. & Corbet, R. 2006, *The Astronomer’s Telegram*, 940, 1
- Leyder, J.-C., Walter, R., Lazos, M., Masetti, N., & Produit, N. 2007, *A&A*, 465, L35
- Maiolino, R., Rieke, G. H., & Rieke, M. J. 1996, *AJ*, 111, 537
- Masetti, N., Mason, E., Morelli, L., et al. 2008, *A&A*, 482, 113
- Molkov, S. V., Cherepashchuk, A. M., Lutovinov, A. A., et al. 2004, *Astronomy Letters*, 30, 534
- Negueruela, I., Smith, D. M., Reig, P., Chaty, S., & Torrejón, J. M. 2006, in *ESA Special Publication, Vol. 604, The X-ray Universe 2005*, ed. A. Wilson, 165–170

- Negueruela, I., Torrejón, J. M., Reig, P., Ribó, M., & Smith, D. M. 2008, in American Institute of Physics Conference Series, Vol. 1010, American Institute of Physics Conference Series, 252–256
- Persson, S. E., Murphy, D. C., Krzeminski, W., Roth, M., & Rieke, M. J. 1998, *AJ*, 116, 2475
- Rahoui, F., Chaty, S., Lagage, P.-O., & Pantin, E. 2008, *A&A*, 484, 801
- Searle, S. C., Prinja, R. K., Massa, D., & Ryans, R. 2008, *A&A*, 481, 777
- Sguera, V., Hill, A. B., Bird, A. J., et al. 2007, *A&A*, 467, 249
- Sidoli, L., Romano, P., Mereghetti, S., et al. 2007, *A&A*, 476, 1307
- Ubertini, P., Lebrun, F., Di Cocco, G., et al. 2003, *A&A*, 411, L131
- Walter, R. & Zurita Heras, J. 2007, *A&A*, 476, 335
- Winkler, C., Courvoisier, T. J.-L., Di Cocco, G., et al. 2003, *A&A*, 411, L1